



**A Meeting to Mark the Sixty-fifth Birthday of Professor
M S P Eastham**

26th-27th July, 2002

**University of Wales
Gregynog Hall
Newtown
Powys**

London
Mathematical Society

Cardiff University
Department of Computer Science

A Meeting to Mark the Sixty-fifth Birthday of Professor M S P Eastham

Gregynog Hall, 26-27th July, 2002

CONTENTS OF THIS BOOKLET

Laudatum	ii
Speakers	iv
Timetable of Talks and Events	v
Abstracts of Presentations	vi
List of Participants	xi
Acknowledgement	xvi

LAUDATUM

B M BROWN AND W D EVANS

Michael Eastham received his early education in Manchester Grammar School and Merton College, Oxford. In 1959, after a successful undergraduate career, he was awarded a Domus Senior Scholarship by Merton College to enable him to pursue postgraduate studies under the guidance of E.C.Titchmarsh. In 1961, in recognition of the excellence of the research for his D.Phil thesis, he was presented with the prestigious Senior Mathematical Scholarship by the University of Oxford. In 1978 he was awarded the Keith Prize and Gold Medal by the Royal Society of Edinburgh for his contribution to classical analysis, and was made a Fellow of the Royal Society of Edinburgh in 1982.

After a year as Junior Research Fellow of Merton College, Michael left Oxford in 1962 to take up an appointment at Reading University. Later he held positions in the universities of Southampton, London (Chelsea College and King's College) where he was Professor of Pure Mathematics 1980-1988, and Bahrain. At present he is an honorary Professorial Fellow of the University of Wales at Cardiff.

He has made a significant contribution to classical analysis, with an impressive output of 4 books and over 100 papers to date. His main area of expertise is the spectral theory of linear ordinary differential equations. Of notable significance is his work on the asymptotics of solutions. Using the technique of repeated diagonalisation with consummate skill and efficacy, he has obtained powerful results on asymptotics in a number of papers which were the basis of his well-known book on the subject. His results have enabled him and his students to make a comprehensive analysis of the deficiency index problem for equations with suitably smooth coefficients. In recent years he has developed his work on asymptotics in a manner suitable for inclusion as an algorithm that can be realised on a computer, and has applied his methods effectively to a problem of considerable current interest, namely, that of the location of Sturm-Liouville resonances. Out of Michael's long and impressive list of publications, the papers listed below are particularly noteworthy, and are good examples of the power of his analysis.

Michael has been a leading figure in the subject for the last forty years. He has lectured on his work in many countries, and his books have become standard texts. He continues to be very active in research, his current main areas of activity being spectral concentration and resonance problems.

SELECTED PUBLICATIONS

1. B. M. Brown and M. S. P. Eastham. Analytic continuation and resonance-free regions for Sturm-Liouville potentials with power decay. *J. Comp. Appl. Math.*, to appear, 2002.

This *Laudatum* will appear in a future edition of the Journal of Computational and Applied Mathematics.

2. B. M. Brown, M. S. P. Eastham, and D. K. R. McCormack. Spectral concentration and rapidly decaying potentials. *J. Comput. Appl. Math.*, 81(2):333–348, 1997.
3. B. M. Brown, M. S. P. Eastham, and D. K. R. McCormack. Resonances and analytic continuation for exponentially decaying Sturm-Liouville potentials. *J. Comput. Appl. Math.*, 116(1):181–193, 2000.
4. M. S. P. Eastham. On the gaps in the spectrum associated with Hill’s equation. *Proc. Amer. Math. Soc.*, 21:643–647, 1969.
5. M. S. P. Eastham. On a limit-point method of Hartman. *Bull. London Math. Soc.*, 4:340–344, 1972.
6. M. S. P. Eastham. Gaps in the essential spectrum of even-order self-adjoint differential operators. *Proc. London Math. Soc. (3)*, (34) :213-230, 1977.
7. M. S. P. Eastham. Asymptotic theory and deficiency indices for differential equations of odd order. *Proc. Roy. Soc. Edinburgh Sect. A*, 90(3-4):263–279, 1981.
8. M. S. P. Eastham. The deficiency index of a second-order differential system. *J. London Math. Soc. (2)*, 23(2):311–320, 1981.
9. M. S. P. Eastham. On the number of solutions of right-definite problems with a convergent Dirichlet integral. *Proc. Roy. Soc. Edinburgh Sect. A*, 91:347–360, 1982.
10. M. S. P. Eastham. The asymptotic solution of linear differential systems. *Mathematika*, 32(1):131–138, 1985.
11. M. S. P. Eastham. The number of resonant states in perturbed harmonic oscillation. *Quart. J. Math. (Oxford) (2)*, 42: 49-55,1991.
12. M. S. P. Eastham. Antibound states and exponentially decaying Sturm-Liouville potentials. *J. London Math. Soc.*, to appear, 2002.
13. M. S. P. Eastham, C. T. Fulton, and S. Pruess. Using the SLEDGE package on Sturm-Liouville problems having nonempty essential spectra. *ACM Trans. Math. Software*, 22:423–446, 1996.
14. M. S. P. Eastham and J. B. McLeod. The existence of eigenvalues embedded in the continuous spectrum of ordinary differential operators. *Proc. Roy. Soc. Edinburgh Sect. A*, 79:25–34, 1977.
15. M. S. P. Eastham and J. B. McLeod. Non-resonance for linear differential systems. *Proc. Roy. Soc. Edinburgh Sect. A*, 111(1-2):103–122, 1989.

SPEAKERS

E B Davies, *Electronic mail:* E.B.Davies@kcl.ac.uk
Kings College, University of London

W D Evans, *Electronic mail:* EvansWD@cardiff.ac.uk
Cardiff University

W N Everitt, *Electronic mail:* w.n.everitt@bham.ac.uk
University of Birmingham

Hubert Kalf, *Electronic mail:* Hubert.Kalf@mathematik.uni-muenchen.de
Universität München

J B McLeod, *Electronic mail:* mcleod@pitt.edu
University of Pittsburgh

TIMETABLE OF TALKS AND EVENTS

FRIDAY, JULY 26TH, 2002

Chair W D Evans

1500 **E B Davies** *Bounds on the eigenvalues of non-self-adjoint Schrödinger operators*

1600 *Tea*

1630-1730 **W N Everitt[†] and Anthippi Poulkou** *First-order linear boundary value problems*

1900 *Reception*

1930 *Dinner*

followed by the *Laudatum* by Prof W N Everitt.

SATURDAY, JULY 27TH, 2002

0800 *Breakfast*

Chair B M Brown

0900 **A A Balinsky and W D Evans[†]** *On the spectral properties of the Brown-Ravenhall operator*

1000 **Hubert Kalf** *On the spectral theory of Dirac operators with a variable mass term*

1100 *Coffee*

1130 **J B McLeod** *Stability of Poiseuille Flow*

1230 *Lunch*

Depart

Where there is more than one author, the speaker is marked thus [†].

ABSTRACTS OF PRESENTATIONS

BOUNDS ON THE EIGENVALUES OF NON-SELF-ADJOINT SCHRÖDINGER OPERATORS

E B DAVIES

E.B.Davies@kcl.ac.uk

Department of Mathematics

Kings College, University of London

The Strand

London WC2R 2LS, UK

This talk describes bounds on the location of the eigenvalues of one-dimensional non-self-adjoint Schrödinger operators, obtained jointly by the speaker and J Nath. The bounds are expressed in terms of L^p norms of the potentials, and are optimal if the potential is L^1 .

ON THE SPECTRAL PROPERTIES OF THE BROWN-RAVENHALL OPERATOR

A A BALINSKY

BalinskyA@cardiff.ac.uk

W D EVANS

EvansWD@cardiff.ac.uk

School of Mathematics

Cardiff University

23 Senghennydd Road

P O Box 926

Cardiff CF24 4YH, UK

The fact that the Dirac is unbounded below creates problems if it is used to describe multi-particle relativistic systems since the resulting operator has a spectrum which covers the whole of the real line. To overcome this difficulty Brown and Ravenhall proposed the following one-particle model. To describe an electron in the field of its nucleus and subject to relativistic effects, the operator of Brown and Ravenhall is

$$(1) \quad \mathbf{B} := \Lambda_+ \left(D_0 - \frac{e^2 Z}{|\cdot|} \right) \Lambda_+.$$

acting in the Hilbert space $\mathcal{H} := \Lambda_+(L^2(\mathbb{R}^3) \otimes \mathbb{C}^4)$. The notation in (1) is as follows

- D_0 is the free Dirac operator

$$D_0 = c\boldsymbol{\alpha} \cdot \frac{\hbar}{i} \nabla + mc^2\beta \equiv \sum_{j=1}^3 c \frac{\hbar}{i} \alpha_j \frac{\partial}{\partial x_j} + mc^2\beta,$$

where $\boldsymbol{\alpha} = (\alpha_1, \alpha_2, \alpha_3)$ and β are the Dirac matrices given by

$$\alpha_j = \begin{pmatrix} 0 & \sigma_j \\ \sigma_j & 0 \end{pmatrix}, \beta = \begin{pmatrix} 1_2 & 0_2 \\ 0_2 & -1_2 \end{pmatrix}$$

with $0_2, 1_2$ the zero and unit 2×2 matrices respectively and σ_j the Pauli matrices

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

- Λ_+ denotes the projection of $L^2(\mathbb{R}^3) \otimes \mathbb{C}^4$ onto the positive spectral subspace of D_0 , that is $\chi_{(0,\infty)}(D_0)$, where $\chi_{(0,\infty)}$ is the characteristic function of $(0, \infty)$. If we set

$$\widehat{f}(\mathbf{p}) \equiv \mathcal{F}(f)(\mathbf{p}) = \left(\frac{1}{2\pi\hbar} \right)^{3/2} \int_{\mathbb{R}^3} e^{-i\mathbf{x}\cdot\mathbf{p}/\hbar} f(\mathbf{x}) d\mathbf{x}$$

for the Fourier transform of f , then it follows that

$$(\Lambda_+ f)^\wedge(\mathbf{p}) = \Lambda_+(\mathbf{p}) \widehat{f}(\mathbf{p}),$$

where

$$(2) \quad \Lambda_+(\mathbf{p}) = \frac{1}{2} + \frac{c\boldsymbol{\alpha} \cdot \mathbf{p} + mc^2\beta}{2e(p)}, \quad e(p) = \sqrt{c^2p^2 + m^2c^4}$$

with $p = |\mathbf{p}|$.

- $2\pi\hbar$ is Planck's constant, c the velocity of light, m the electron mass, $-e$ the electron charge, and Z the nuclear charge.

The lecture will discuss spectral properties of operators $b_{l,s}$ appearing in the partial wave decomposition of \mathbf{B} : the indices l, s , denote the angular momentum channel and spin respectively. The following topics will be covered: the value of the critical charge $Z_c(l, s)$ which yields the positivity of $b_{l,s}$, the charge range for essential self-adjointness, and the charge range for the absence of embedded eigenvalues.

FIRST-ORDER LINEAR BOUNDARY VALUE PROBLEMS

W N EVERITT

w.n.everitt@bham.ac.uk

School of Mathematics and Statistics

University of Birmingham

Edgbaston

Birmingham B15 2TT, UK

ANTHIPPI POULKOU

apoulkou@cc.uoa.gr

Department of Mathematics

University of Athens

Panepistemiopolis

Athens 157 84, Greece

1. ABSTRACT

This lecture reports on joint work with Anthippi Poulkou, Department of Mathematics, University of Athens.

The general Lagrange symmetric first-order differential equation with Lebesgue integrable coefficients, on the open interval (a, b) of the real line \mathbb{R} , has the form, defining the differential expression $M[\cdot]$,

$$M[y](x) := i\rho(x)y'(x) + \frac{1}{2}i\rho'(x)y(x) + q(x)y(x) = \lambda w(x)y(x) \text{ for all } x \in (a, b)$$

where $\lambda \in \mathbb{C}$ is the complex spectral parameter. Here the coefficients ρ, q, w satisfy the conditions

- (i) $\rho, q, w : (a, b) \rightarrow \mathbb{R}$
- (ii) $\rho \in AC_{\text{loc}}(a, b)$ and $\rho(x) > 0$ for all $x \in (a, b)$
- (iii) $q, w \in L^1_{\text{loc}}(a, b)$
- (iv) $w(x) > 0$ for almost all $x \in (a, b)$.

The right-definite spectral analysis for this differential equation takes place in the Hilbert function space $L^2((a, b); w)$ with norm and inner-product

$$\|f\|_w^2 := \int_I w|f|^2 \text{ and } (f, g)_w := \int_a^b w(x)f(x)\bar{g}(x) dx.$$

A necessary and sufficient condition to ensure that the differential expression $M[\cdot]$ generates a maximal operator in $L^2((a, b); w)$ with equal deficiency indices $d^\pm = 1$ whose self-adjoint restrictions have discrete spectra, is

$$\int_a^b \frac{w(x)}{\rho(x)} dx < +\infty.$$

With this condition satisfied the GKN boundary condition method can be applied to give symmetric boundary value problems with the following properties:

Theorem 1.1. *Let T be a self-adjoint restriction of the maximal operator generated by $M[\cdot]$; then T has the following spectral properties:*

- (i) *The spectrum $\sigma(T)$ of T in $L^2((a, b); w)$ is simple and discrete.*
- (ii) *The spectrum $\sigma(T)$ is unbounded above and below on $\mathbb{R} \subset \mathbb{C}$, and so may be denoted by, here $\mathbb{Z} = \{\dots - 2, -1, 0, 1, 2, \dots\}$,*

$$\sigma(T) = \{\lambda_n \in \mathbb{R} : n \in \mathbb{Z}\}$$

with

$$\lambda_n < \lambda_{n+1} \text{ for all } n \in \mathbb{Z}, \text{ and } \lim_{n \rightarrow \pm\infty} \lambda_n = \pm\infty.$$

- (iii) *There exists a positive number $k > 0$, with*

$$k = 2\pi \left(\int_a^b \frac{w(x)}{\rho(x)} dx \right)^{-1},$$

such that

$$\lambda_{n+1} - \lambda_n = k \text{ for all } n \in \mathbb{Z}.$$

- (iv) *There exists an entire (integral) function $\varphi : \mathbb{C} \rightarrow \mathbb{C}$, generated by the boundary value problem, with the properties*

- (i) $\varphi(\lambda) = 0$ if and only if $\lambda \in \{\lambda_n : n \in \mathbb{Z}\}$
- (ii) $\varphi'(\lambda_n) \neq 0$ for all $n \in \mathbb{Z}$.

2. KRAMER ANALYTIC KERNELS

The boundary value problems discussed in Section 1 generate Kramer analytic kernels in the Hilbert space $L^2((a, b); w)$.

Acknowledgement The authors are indebted to the Professors Michael Plum and Hubert Kalf for technical help in the preparation of the manuscript and for correcting errors in the first draft of the paper.

References

- [1] W.N. Everitt and L. Markus. ‘The Glazman-Krein-Naimark theorem for ordinary differential operators.’ *Operator Theory: Advances and Applications* **98** (1997), 118-130.
- [2] W.N. Everitt, G. Nasri-Roudsari and J. Rehberg. ‘A note on the analytic form of the Kramer sampling theorem.’ *Results in Mathematics*. **34** (1998), 310-319.
- [3] W.N. Everitt and Anthippi Poulkou. ‘Kramer analytic kernels and first-order boundary value problems.’ *Jour. Computational Appl. Math.* (To appear.)

ON THE SPECTRAL THEORY OF DIRAC OPERATORS WITH A VARIABLE MASS TERM

HUBERT KALF

Hubert.Kalf@mathematik.uni-muenchen.de

Mathematisches Institut

Universität München

Theresienstr. 39

D-80333 München, Germany

The spectrum of the Dirac operator is purely discrete when the mass term “dominates” the potential (O. Yamada). In the opposite case one expects the spectrum to be purely absolutely continuous. This was proved when both the mass term and the potential are spherically symmetric (K. M. Schmidt, O. Yamada). Using virial techniques, a theorem is presented which establishes at least absence of eigenvalues when mass term and potential are not necessarily rotationally symmetric. This is joint work with T. Okaji (Kyoto) and O. Yamada (Kusatsu).

STABILITY OF POISEUILLE FLOW

J B MCLEOD

mcleod@pitt.edu

Department of Mathematics

301 Thackeray Hall

University of Pittsburgh

Pittsburgh PA, 15260, USA

Poiseuille flow is two-dimensional flow in a straight pipe. The question of the stability of the steady flow, particularly with a parabolic velocity profile, is a long-standing one that has received intense treatment numerically, but very little analytically. The talk will examine what we can prove about this problem.

LIST OF PARTICIPANTS

Professor A A Abramov, *Electronic mail:* alalabr@ccas.ru

Department of Computational Methods, Dorodnicyn Computing Centre of the Russian Academy of Sciences, Vavilov St. 40, 119991 Moscow GSP-1, Russia

Dr A A Balinsky, *Electronic mail:* BalinskyA@cardiff.ac.uk

School of Mathematics, Cardiff University, 23 Senghennydd Road, P O Box 926, Cardiff CF24 4YH, UK

Dr H Behnke, *Electronic mail:* mahb@sinfonix.rz.tu-clausthal.de

Institut für Mathematik, Technische Universität Clausthal, Erzstraße 1, 38678 Clausthal-Zellerfeld, Germany

Dr Christer Bennewitz, *Electronic mail:* christer.bennewitz@math.lu.se

Department of Mathematics, Lund University, Box 118, SE-221 00 Lund, Sweden

Professor Paul Binding, *Electronic mail:* binding@math.ucalgary.ca

The Department of Mathematics and Statistics, University of Calgary, 2500 University Drive, Calgary, Alberta, Canada, T2N 1N4

Dipl.-Math B Breuer, *Electronic mail:* bodo@ma1euler.mathematik.uni-karlsruhe.de

Mathematisches Institut I, Universität Karlsruhe, D-76128 Karlsruhe, Germany

Dr B M Brown, *Electronic mail:* Malcolm.Brown@cs.cardiff.ac.uk

Department of Computer Science, Cardiff University, P O Box 916, Cardiff CF24 3XF, UK

Professor Patrick J Browne, *Electronic mail:* browne@admin.usask.ca

Department of Mathematics and Statistics, University of Saskatchewan, Saskatoon, Saskatchewan S7N 5E6, Canada

Professor E B Davies FRS, *Electronic mail:* E.B.Davies@kcl.ac.uk

Department of Mathematics, Kings College, University of London, The Strand, London WC2R 2LS, UK

Professor Michael S P Eastham, *Electronic mail:* mandh@chesilhay.fsnet.co.uk

Department of Computer Science, Cardiff University, PO Box 916, Cardiff CF24 3XF, UK

Professor W D Evans, *Electronic mail:* EvansWD@cardiff.ac.uk

School of Mathematics, Cardiff University, 23 Senghennydd Road, P O Box 926, Cardiff CF24 4YH, UK

Professor W N Everitt, *Electronic mail:* w.n.everitt@bham.ac.uk

School of Mathematics and Statistics, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

Dr D J Gilbert, *Electronic mail:* dgilbert@maths.kst.dit.ie

School of Mathematical Sciences, Dublin Institute of Technology, Kevin Street, Dublin 8, Ireland

Professor Leon Greenberg, *Electronic mail:* lng@math.umd.edu

Department of Mathematics, University of Maryland, College Park, Maryland MD 20742 , USA

Dr G Gudmundsdottir, *Electronic mail:* Gudmundsdottir@math.lth.se

Department of Mathematics, Lund University, Box 118, SE-221 00 Lund, Sweden

Professor Dr Rainer Hempel, *Electronic mail:* hempel@rzserv6i.rz.tu-bs.de

Institut für Analysis, TU Braunschweig, Germany

Professor D Hinton, *Electronic mail:* hinton@math.utk.edu

Mathematics Department, , University of Tennessee,, Knoxville, , TN 37996-1300, USA

Professor Dr Andreas M Hinz, *Electronic mail:* hinz@rz.mathematik.uni-muenchen.de

Zentrum Mathematik, Technische Universität München, D-80290 München, Germany

Dr Michael Hitrik, *Electronic mail:* hitrik@math.berkeley.edu

Department of Mathematics , University of California at Berkeley , Berkeley, CA 94720-3840 , USA

M Jais, *Electronic mail:* mathiasjais@gmx.net

Zentrum Mathematik, Technische Universität München, D-80290 München, Germany

Department of Computer Science, Cardiff University, P O Box 916, Cardiff CF24 3XF, UK

Professor Dr Hubert Kalf, *Electronic mail:* Hubert.Kalf@mathematik.uni-muenchen.de

Mathematisches Institut, Universität München, Theresienstr. 39, D-80333 München, Germany

Professor Ian Knowles, *Electronic mail:* iwk@vorteb.math.uab.edu

Department of Mathematics, University of Alabama at Birmingham, University Station, Birmingham, AL 35294, USA

- O A Krus**, *Electronic mail*: tchebolga@leibniz.iimas.unam.mx
 IIMAS, UNAM Apdo Postal 20-726, Admon No 20. Deleg. Alvaro Obregon, 01000 Mexico DF, Mexico
- Dr P Kurasov**, *Electronic mail*: kurasov@maths.lth.se
 Centre for Mathematical Sciences, Lund Institute of Technology, Lund University, Box 118, SE-221 00 Lund, Sweden
- Professor Dr Heinz Langer**, *Electronic mail*: hlanger@mail.zserv.tuwien.ac.at
 Institute for Analysis and Technical Mathematics, Technical University of Vienna, Wiedner Hauptstr. 8-10, A-1040 Vienna, Austria
- Dr Matthias Langer**, *Electronic mail*: mlanger@math.uni-bremen.de
 FB 3 - Mathematik, Universität Bremen, Bibliothekstrasse 1, D-28359 Bremen, Germany
- Professor Roger T Lewis**, *Electronic mail*: rlewis@nsf.gov
 Division of Mathematical Sciences, Suite 1025, National Science Foundation, 4201 Wilson Blvd., Arlington, VA 22230, USA
 Department of Mathematics, University of Alabama at Birmingham, University Station, Birmingham AL 35294, USA
- Professor Juan J Manfredi**, *Electronic mail*: manfredi@pitt.edu
 Department of Mathematics, 301 Thackeray Hall, University of Pittsburgh, Pittsburgh PA, 15260, USA
 Division of Mathematical Sciences, Suite 1025, National Science Foundation, 4201 Wilson Blvd., Arlington, VA 22230, USA
- Dr Marco Marletta**, *Electronic mail*: mm7@le.ac.uk
 Department of Mathematics and Computer Science, University of Leicester, University Road, Leicester LE1 7RH, UK
- Professor J B McLeod FRS**, *Electronic mail*: mcLeod@pitt.edu
 Department of Mathematics, 301 Thackeray Hall, University of Pittsburgh, Pittsburgh PA, 15260, USA
- Professor Sergey Naboko**, *Electronic mail*: naboko@felix.math.uab.edu
 Department of Mathematical Physics, Institute for Physics, St Petersburg State University, Ulyanovskaya 1, St Petersburg 198904, Russia
- Professor B S Pavlov**, *Electronic mail*: pavlov@aitken.scitec.auckland.ac.nz
 Department of Mathematical Physics, Institute for Physics, St Petersburg State University, Ulyanovskaya 1, St Petersburg 198904, Russia
 Department of Mathematics, University of Auckland, Private Bag 92019, Auckland, New Zealand

Professor D B Pearson, *Electronic mail*: d.b.pearson@hull.ac.uk

Department of Mathematics, University of Hull, Hull HU6 7RX, UK

Professor Dr M Plum, *Electronic mail*: michael.plum@math.uni-karlsruhe.de

Mathematisches Institut I, Universität Karlsruhe, D-76128 Karlsruhe, Germany

Dipl.-Math R Pohler, *Electronic mail*: rudi@ma1euler.mathematik.uni-karlsruhe.de

Mathematisches Institut I, Universität Karlsruhe, D-76128 Karlsruhe, Germany

Dr A Pushnitski, *Electronic mail*: A.B.Pushnitski@lboro.ac.uk

Department of Mathematical Sciences, Loughborough University, Loughborough, Leicestershire LE11 3TU, UK

Dr Wolfgang Reichel, *Electronic mail*: reichel@math.unibas.ch

Mathematisches Institut, Universität Basel, Rheinsprung 21, CH-4051 Basel, Switzerland

Dr Norbert Röhrli, *Electronic mail*: ngr@vorteb.math.uab.edu

Department of Mathematics, The University of Alabama at Birmingham, Campbell Hall, Birmingham, AL 35294-1170, USA

Dipl.-Math J Rohe, *Electronic mail*: uafx@rz.uni-karlsruhe.de

Mathematisches Institut I, Universität Karlsruhe, D-76128 Karlsruhe, Germany

Dr R Romanov, *Electronic mail*: R.Romanov@cs.cardiff.ac.uk

Department of Computer Science, Cardiff University, PO Box 916, Cardiff, CF24 3XF, UK

V S Samko, *Electronic mail*: V.Samko@cs.cardiff.ac.uk

Department of Computer Science, Cardiff University, PO Box 916, Cardiff, CF24 3XF, UK

Dipl.-Math M Schäfer, *Electronic mail*: mschaefer@uni-duisburg.de

Gerhard-Mercator-Universität, Fakultät 4 (Mathematik), D-47048 Duisburg, Germany

Dr Karl Michael Schmidt, *Electronic mail*: SchmidtKM@cardiff.ac.uk

School of Mathematics, Cardiff University, Senghennydd Road, Cardiff CF24 4YH, UK

Professor A A Shkalikov, *Electronic mail*: ashkalikov@yahoo.com

Department of Mechanics and Mathematics, Moscow Lomonosov State University, Moscow, 198 899, Russia

Ian Sorell, *Electronic mail*: i.sorell@lboro.ac.uk

Department of Mathematical Sciences, Loughborough University, Loughborough, Leicestershire LE11 3TU, UK

Professor Gunter Stoltz, *Electronic mail*: stolz@vorteb.math.uab.edu

Department of Mathematics, University of Alabama at Birmingham, University Station, Birmingham AL 35294, USA

Professor Dr Christiane Tretter, *Electronic mail*: ctretter@math.uni-bremen.de

FB 3 - Mathematik, Universität Bremen, Bibliothekstrasse 1, D-28359 Bremen, Germany

Dr A Watson, *Electronic mail*: WatsonA1@cardiff.ac.uk

School of Engineering, Cardiff University, PO Box 925, Cardiff, CF24 0YF, UK

Professor R Weikard, *Electronic mail*: rudi@math.uab.edu

Department of Mathematics, University of Alabama at Birmingham, University Station, Birmingham AL 35294, USA

Dr Robin Williams, *Electronic mail*: Robin.Williams@astro.cf.ac.uk

Department of Physics and Astronomy, Cardiff University, PO Box 913, Cardiff, CF24 3YB, UK

Professor A Zettl, *Electronic mail*: zettl@math.niu.edu

Department of Mathematical Sciences, Northern Illinois University, Dekalb, Illinois 60115, USA

ACKNOWLEDGEMENT

This meeting is organised by B M Brown and W D Evans and is in part sponsored by the Department of Computer Science, Cardiff University and the London Mathematical Society.